

Diffractive Jet Production in DIS

—

Probing the Structure of Colour Singlet Exchange

Frank-Peter Schilling / DESY

<http://www.desy.de/~fpschill>

H1 Collaboration





Fermilab and Madison

May 2001

- Introduction
- Diffractive DIS and $F_2^{D(3)}$
- Diffractive jet production
- Results
- Summary and conclusions


HERA and the H1 Detector




H1, ZEUS: $e p$ collisions at
 $\sqrt{s} = 320\text{ GeV}$

HERA-B: p -beam on fixed target:
 CP violation in $B^0 \bar{B}^0$

HERMES: e -beam on polarized target:
 Spin structure



Deep Inelastic Scattering (DIS) at HERA





$$Q^2 = -q^2 = (k - k')^2$$

Photon virtuality,
"Resolution power"

$$x = \frac{-q^2}{2P \cdot q} \quad (0 < x < 1)$$

Parton momentum fraction in p


- Highly virtual point-like photon γ^* in DIS at HERA probes proton structure with unprecedented resolution




- Scattering off coloured object:
→ p breaks up ("proton remnant")

Large Rapidity Gap (LRG) Events

- 10% of DIS events for exhibit large gap without hadronic activity in outgoing \mathbf{p} region



- γ^* scatters off colorless state in \mathbf{p} ("Pomeron")
- \mathbf{p} (or low-mass excitation) escapes through beampipe



$$t = (p - p')^2 : \text{(momentum transfer)}^2 \text{ at } \mathbf{p} \text{ vertex}$$

$$M_X, M_Y : \text{Masses of } \mathbf{X} \text{ and } \mathbf{Y}$$

$$x_{IP} = \frac{q \cdot (p - Y)}{q \cdot p} = \frac{Q^2 + M_X^2 - t}{Q^2 + W^2 - M_p^2}$$


→ long. momentum fraction transferred from \mathbf{p} to exchange

$$\beta = \frac{-q^2}{q \cdot (p - Y)} = \frac{Q^2}{Q^2 + M_X^2 - t}$$

→ fraction of exchange momentum carried by \mathbf{q} coupling to γ

Reminder: The “Pomeron”

- Introduced as pseudo-particle to parameterize elastic high energy scattering at small momentum transfers:



- Pomeron trajectory:

$$\alpha(t) = \alpha(0) + \alpha' t = 1.08 + 0.25 t$$
- Differential and total cross section:


$$\frac{d\sigma}{dt} \sim \frac{1}{s^2} |T(s, t)|^2 = f(t) \left(\frac{s}{s_0}\right)^{2\alpha(t)-2}$$

$$\sigma_{tot} \sim \frac{1}{s} \text{Im}(T(s, t))|_{(t=0)} = s^{\alpha(0)-1}$$


⇒ Today: Understand colour singlet exchange in terms of QCD (quark and gluon dynamics)!

Selection of LRG Events


1. Tagging of p with “Roman Pots” (measure t , small stat.):




2. Large Rapidity Gap Requirement (integr. over M_Y, t):



3. Analysis of final state M_X system (integr. over M_Y, t):



Colour Singlet Exchange Processes in $\gamma^* p$ Interactions




- Photon γ^* can either fluctuate into vector meson or dissociate into high-mass system X
- Proton p either stays intact (elastic scattering) or dissociates into low-mass baryonic system Y

Diffractive DIS: Probing $\textbf{\textit{IP}}$ Structure

Inclusive DIS: Structure function $F_2(x, Q^2)$:

$$\frac{d^2\sigma(\text{incl.})}{dx dQ^2} = \frac{4\pi\alpha^2}{xQ^4} \left(1 - y + \frac{y^2}{2}\right) F_2(x, Q^2)$$



Diffractive DIS: Diffractive structure function F_2^D :

$$\frac{d^5\sigma(ep \rightarrow eXY)}{dx_{\text{IP}} dt dM_Y d\beta dQ^2} = \frac{4\pi\alpha^2}{\beta Q^4} \left(1 - y + \frac{y^2}{2}\right) \times F_2^{D(5)}(x_{\text{IP}}, t, M_Y, \beta, Q^2)$$

If Y is not measured, integrate over M_Y, t :


$$F_2^{D(3)}(\beta, Q^2, x_{\text{IP}}) = \int_{m_P}^{M_Y, \text{max}} dM_Y \int_{t_2}^{t_1} dt F_2^{D(5)}$$

\Rightarrow Large gap between X and $Y \Leftrightarrow M_X, M_Y, x_{\text{IP}}$ small

The Diffractive Structure Function $F_2^{D(3)}$

H1 Measurement of $F_2^{D(3)}(x_{IP}, \beta, Q^2)$:

$$x_{IP} F_2^D \quad \beta \rightarrow$$



Diffractive Parton Distributions

- Inclusive DIS factorization theorem:


$$F_2(x, Q^2) \sim C_i \otimes p_i \quad (+\text{higher twist})$$

- Diffractive DIS [proof by J. Collins in 1998]:

$$F_2(x, Q^2, x_{IP}, t) \sim C_i \otimes p_i^D \quad (+\text{higher twist})$$

- valid at fixed x_{IP}, t
- p_i^D : ‘conditional probabilities’, obey DGLAP evolution
- determine p_i^D in inclusive diffr. scattering, then predict exclusive processes
- same C_i as in inclusive DIS

- Additional assumption: factorizing x_{IP} dependence
[‘Regge factorization’]




$$F_2^{D(3)}(x_{IP}, \beta, Q^2) = f_{IP/p}(x_{IP}) \times F_2^{IP}(\beta, Q^2)$$

Diffractive PDF's from $F_2^{D(3)}$


- Observation of (positive!) scaling violations:

$F_2^D \quad (\beta = 0.4)$




[flat/rising with Q^2]

$F_2 \quad (x = 0.4)$



[steeply falling with Q^2]

- Strongly suggestive of g dominated exchange !
- DGLAP QCD analysis of F_2^D yields diffractive PDF's:



Gluons
»
Quarks


$[g^D$ indirectly
determined from
scal. viol.
→ large uncertainty]

Diffractive Dijet Production

Motivation:

- Direct sensitivity to g^D through $\mathcal{O}(\alpha_s)$ process (boson gluon fusion):
- Jet P_T provides second hard scale

Kinematics:



M_{12}


- Invariant mass of two leading jets

$$z_{IP}^{(jets)} \approx \frac{Q^2 + M_{12}^2}{Q^2 + M_X^2}$$

- Momentum fraction of exch. entering hard scattering

Proton Rest Frame Picture

- Photon fluctuates into $q\bar{q}$ or $q\bar{q}g$ state long before interaction, then scatters elastically off proton at rest




$q\bar{q}$ \Leftrightarrow diffr. q scattering (quark parton model)

$q\bar{q}g$ \Leftrightarrow diffr. g scattering (boson gluon fusion)

- For large diffractive masses M_X or high p_T final states, $q\bar{q}g$ configurations are expected to dominate

Event Selection



- DIS Signature: $4 < Q^2 < 80$ GeV; $0.1 < y < 0.7$
Scattered electron e'
- Diffractive Signature: $x_P < 0.05$; M_Y, t small
Rapidity gap in outgoing p' direction ($3.2 < \eta < 7.5$)
- 2-Jet Signature: $N_{\text{Jet}} \geq 2$; $p_T > 4$ GeV
CDF cone jet-algorithm in γp CMS

$$\mathcal{L} = 18.0 \text{ pb}^{-1} \quad N_{\text{2-Jet}} = 2.500 \quad N_{\text{3-Jet}} = 130$$

Cross Section Measurement

- Correction for detector effects to ‘stable’ particle level
- Correction for QED radiative effects at lepton vertex
- Bin-by-bin correction procedure



Kinematic Range of Hadron Level Cross Sections	
$4 < Q^2 < 80 \text{ GeV}^2$	
$0.1 < y < 0.7$	
$x_{IP} < 0.05$	
$M_Y < 1.6 \text{ GeV}$	
$ t < 1.0 \text{ GeV}^2$	
$N_{\text{jets}} \geq 2$ or $N_{\text{jets}} = 3$	
$p_{T,jet}^* > 4 \text{ GeV}$	
$-3 < \eta_{jet}^* < 0$	

Main systematic uncertainties:

- Hadronic energy scales of calorimeters
- Trigger efficiency
- Model dependence of correction
- Scattered electron measurement

⇒ Systematic error dominates (15 – 30%)

Results: General Properties of Dijet Events




- Significant energy not contained in dijets, some preference for \textbf{IP} hemisphere
 - $M_{12} \ll M_X$ typically
- ⇒ exclusive 2-jets ($q\bar{q}$) just small part of cross section !

QCD Factorization at Work

Predict diffr. dijet cross sections with PDF's obtained from inclusive $F_2^{D(3)}$ measurement:


[resolved γ^* component included]



⇒ Consistent with QCD factorization in diffractive DIS

Diffractive Gluon Distribution

Dijets directly constrain shape and normalization of g^D :



[res. γ^* , $\mathbf{I}\mathbf{R}$ and quark contributions small]


- H1 fit 2: very good agreement with data
- H1 fit 3: overshoots at high z_{IP}
- ACTW-D (Alvero, Collins, Terron, Whitmore fits): too high

⇒ Strong support for fully factorizable diffr. PDF's in DIS which are gluon-dominated with momentum distr. flat in z

Proton rest frame picture: $q\bar{q}g \gg q\bar{q}$ states

Features of Diffractive PDF's


H1 Diffractive Dijets



- Data consistent with DGLAP evolution of PDF's with factorization scale $\mu^2 = Q^2 + p_T^2$
- Also compatible with factorization of x_{IP} dependence $[f_{IP/P}(x_{IP}) \times p_i^D(z, \mu^2)]$
No visible variation of $\alpha_{IP}(0)$ with z_{IP}

Energy Dependence $\alpha_{IP}(0)$

- Shape of x_{IP} distribution sensitive to energy dependence of cross section:



Parameterization used (Regge motivated):

$$f_{IP/P}(x_{IP}, t) \sim \left(\frac{1}{x_{IP}}\right)^{2\alpha_{IP}(t)-1} e^{Bt}$$

$$\alpha_{IP}(t) = \alpha_{IP}(0) + \alpha'_{IP} t \quad [B = 4.6 \text{ GeV}^{-2}, \alpha'_{IP} = 0.26 \text{ GeV}^{-2}]$$

Fit Result:

$$\alpha_{IP}(0) = 1.17^{+0.03}_{-0.03} \text{ (stat.)}^{+0.06}_{-0.06} \text{ (syst.)}^{+0.03}_{-0.07} \text{ (model)}$$

⇒ Consistent with H1- $F_2^{D(3)}$ [Q^2 similar]

- β distribution: Jets are small β , compared with F_2^D


Resolved Virtual Photon Contribution

Since $Q^2 > 4 \text{ GeV}^2$, $p_T^2 > 16 \text{ GeV}^2$

\Rightarrow Jets can “resolve” virtual photon

[expected from inclusive dijet production]

H1 Diffractive Dijets




$$x_{\gamma}^{(jets)} = \frac{(E - p_Z)_{jets}}{(E - p_Z)_X} \quad E_{rem} \text{ in } \gamma^* \text{ hemisphere}$$

Resolved γ^* contribution according to
“SaS-2D” parameterization [Schuler,Sjöstrand]


- $x_{\gamma}^{(jets)}$ cross section:
Improvement at low $x_{\gamma}^{(jets)}$ if resolved contribution is added
- Corresponding improvement at high E_{rem}^{γ}

Soft Colour Neutralization

- Soft Colour Interactions (SCI)
[Edin, Ingelman, Rathsman]




- Standard DIS, add. soft colour rearrangemens of final state partons may result in rapidity gap
- Two versions:
 - Original version: One additional parameter (rearrangement prob), tuned to F_2^D
 - Improved version: “Generalized area law”; improved description of F_2^D at low Q^2
- Semi-classical model
[Buchmüller, Gehrmann, Hebecker]



- $q\bar{q}, q\bar{q}g$ states scatter elastically off large p , treated as superposition of uncorrelated colour fields

Results: Soft Colour Neutralization




- SCI (orig.) and semi-classical model give reasonable shapes, but underestimate 2-jet cross section by factor 2
- SCI (area law) reproduces normalization, but fails in shapes

⇒ Sensitivity to differences between models which all (have been tuned to) describe $F_2^{D(3)}$!

Colour Dipole / 2-gluon Exchange Models

- Simplest parton level realization of colour singlet exchange: two gluons with cancelling colour charges



- 'Dipole picture': $q\bar{q}, q\bar{q}g$ configurations form (effective) colour dipole scattering off proton

Diffractive cross section:

$$\left. \frac{d\sigma_{T,L}^{\gamma^* p}}{dt} \right|_{t=0} \sim \int d^2 r \int_0^1 d\alpha |\Psi_{T,L}(\alpha, r)|^2 \hat{\sigma}(r^2, x, \dots)$$


Dipole cross section may be expressed as:

$$\hat{\sigma}(x, r) \sim \int \frac{d^2 l_t}{l_t^2} \left[1 - e^{i r \cdot l} \right] \alpha_s(l_t^2) \mathcal{F}(x, l_t^2)$$

Where $\mathcal{F}(x, l_t^2)$ is un-integrated gluon density in proton

Colour Dipole / 2-gluon Exchange Models (II)

- Small P_T , large size dipoles:
similar to soft hadron hadron scattering
- High P_T , small size dipoles:
perturbation theory may be applicable



Saturation Model:
[Golec-Biernat, Wüsthoff]


- Ansatz for σ_{Dipole} which interpolates between pert. ($\sim 1/Q^2$) and non-pert. ($\sim \text{const.}$) parts of inclusive $F_2(x, Q^2)$
- parameters fixed by fit to $F_2(x, Q^2)$, σ^D then predicted
- Strong p_T ordering assumed: $p_{T,g} \ll p_{T,q}$

2-gluon exchange calculation by Bartels et al. (BJLW):
[Bartels, Jung, Lotter, Kyrieleis, Wüsthoff]

- Perturbative calculation in low- β , low- x_{IP} limit
- For $q\bar{q}g$ require high p_T of all 3 partons (only for jets!)
- non- p_T ordered configurations included,
need cut-off for $p_{T,g}$

Results: 2-Gluon Exchange

$x_{IP} < 0.01 \Rightarrow$ avoid IR exch.; p PDF's g -dominated




- tiny $q\bar{q}$ contribution
- BJLW \sim OK if $p_{T,g} > 1.5 \text{ GeV}$
- Saturation Model too low
- $p_{T,rem}^{(IP)}$ not able to discriminate ;-(

3-Jet Production

Features:

- Limited statistics: 130 3-jets for $\mathcal{L} = 18.0 \text{ pb}^{-1}$
- Kinematically forced to $x_{IP} > 0.01$



- Data above LO QCD prediction based on diffr. PDF's if MEPS is used for higher order approximation
- CDM does better job


[Difference MEPS/CDM much smaller for dijets]

- 2-gluon exchange (BJLW) low

Comparison with Tevatron Results

CDF measurement of diffractive dijet production
with leading anti-proton in $p\bar{p}$ collisions:

Effective diffr. structure function $\bar{F}_{JJ}^D(\beta)$



- Prediction based on diffractive PDF's extracted at HERA one order of magnitude above measured cross section!

⇒ Serious breaking of factorization!

Important to understand to get unified picture!


Summary

- Diffr. dijets tightly constrain diffractive gluon distribution g^D (shape and norm.), in contrast to $F_2^{D(3)}$ measurements
- Data favour diffr. PDF's, evolving with DGLAP, strongly dominated by gluons with momentum distribution rel. flat in z ("H1 fit 2")
- Consistent picture from $F_2^{D(3)}$ and jet measurements : Concept of factorizing diffr. PDF's in DIS [Collins] works.
- Consistent with factorizing x_{IP} dependence with $\alpha_{IP}(0) = 1.17$ ("Regge factorization")
- In P rest frame: $q\bar{q}g \gg q\bar{q}$ configurations
- SCI and Semiclassical models not yet able to simultaneously give correct shape and normalizations of jet cross sections
- Improved models calculations based on 2-gluon exchange can describe part of dijet cross section

Conclusion

Major step forward in understanding diffraction:

- Diffractive jet production is a powerful tool to illuminate QCD structure of colour singlet exchange, in particular the role of gluons !
- Sensitivity to discriminate between models which all describe inclusive diffraction (F_2^D) !
- 2-gluon exchange calculations in agreement with diffractive DIS data for the first time !



Results presented here are published by the H1 collaboration:

H1 Collaboration, C. Adloff *et al.*, "Diffractive Jet Production in Deep-Inelastic e^+p Collisions at HERA", DESY 00-174, hep-ex/0012051, acc. by Eur. Phys. J. **C**